

Maga,

Sirup za grlo, uputstvo za upotrebu

Svojstva

Maga sirup za grlo je umirujući sirup za terapiju iritacije grla povezane sa suvim kašljem kod beba i dece. Sadrži glicerol koji ostavlja zaštitni film na sluznici grla i tako postiže antitusivni i umirujući efekat.

Kada se koristi?

Sirup smiruje iritaciju grla povezanu sa suvim kašljem. Štiti od spoljnih faktora kao što su suv vazduh, prašina, dim cigareta i slično.

Uputstvo za upotrebu

Koristite priloženu pipetu za doziranje.

Ubacite pipetu u bočicu sa sirupom, povucite klip pipete i izvucite sirup sve do crne linije. Ta količina sirupa u pipeti je jedna doza od 5 ml.

Vrh pipete stavite u bebina usta. Nežno pritisakajte klip pipete tako da sirup polako ulazi u usta, da bi beba mogla da ga progruta.

Doziranje

Bebe od 6 do 12 meseci: 1 x 5 ml, maksimalno 3 do 4 puta dnevno.

Deca od 1 do 6 godina: 2 x 5 ml, maksimalno 3 do 4 puta dnevno.

Deca od 6 do 12 godina: 1 supena kašika (15 ml), maksimalno 3 do 4 puta dnevno.

Odrasli i deca iznad 12 godina: 1 supena kašika (15 ml), maksimalno 4 do 6 puta dnevno.

Nemojte uzimati veće doze od ovde preporučenih.

Ukoliko zaboravite da uzmete dozu sirupa, nemojte uzimati duplu dozu.

Pipetu morate oprati vodom posle svake upotrebe.

Nemojte prati pipetu u mašini za pranje sudova, nemojte je stavljati u sterilizator za flašice, nemojte je stavljati u mikrotalasnu pećnicu.

Upozorenja

Ne koristite sirup:

- Kod beba mlađih od 6 meseci
- Kod osoba alergičnih na neku od komponenti sirupa
- Kod osoba sa naslednom intolerancijom na fruktozu

Preporučuje se da se posavetujete sa svojim doktorom kod davanja sirupa bebam u starosti od 6 do 12 meseci.

Takođe se posavetujete sa doktorom ukoliko suvi kašlj prate neki teži simptomi (otežano disanje, povišena telesna temperatura koja traje duže od 3 dana, ponavljano povraćanje, problemi sa hranjenjem, dijareja, promene u ponašanju ili problemi sa spavanjem).

Izbegnite da sirup dođe u kontakt sa očima i nosem.

Ukoliko ipak dođe do kontakta sa nosem, očima ili ušima bebe/deteta, neophodno je da odmah isperete sirup, po mogućnosti sa fiziološkim rastvorom.

Držite pipetu van domaćaja dece.

Sastav

Glicerol, prečišćena voda, sirup maltitol, monohidrat limunske kiseline, ukus jagode, kalijum sorbat, natrijum benzoat

Rok trajanja

Čuvati na temperaturi ispod 25°C.

ČUVATI VAN DOMAŠAJA I VIDOKRUGA DECE.

Ne koristiti Maga sirup za grlo posle isteka roka upotrebe naznačenog na kutiji i unutrašnjem pakovanju.

Nakon prvog otvaranja boćice, sirup se mora iskoristiti u roku od 3 (tri) meseca.

Pakovanje

Boćica od 180 ml i pipeta za doziranje od 5 ml.

Datum dobijanja CE znaka: 2016

Datum poslednje revizije uputstva za upotrebu: april 2020



Proizvođač:

Laboratoria Qualipharm, Rijksweg 9, 2880 Bornem, Belgija



Proizvođač (pipeta):

Bormioli Pharma S.p.A. Via Tortona 48, 27055 Rivanazzano (PV), Italija



Ovlašćeni predstavnik, uvoznik i distributer:

EUPHARM phc d.o.o. Bilećka 37,

11000 Beograd, Republika Srbija

www.eupharm.com

www.drsmart.rs

Broj rešenja o registraciji: 515-02-03192-22-003

Kataloški broj: 62950041

Maga, Syrup, information leaflet

Properties

Maga syrup is a soothing syrup for the treatment of an irritated throat associated with a dry cough in babies and children. It contains glycerol, which leaves a protective film in the throat that has an antitussive and soothing effect.

When to use

It calms the irritation of the throat associated with a dry cough. It protects against environmental factors (such as dry air, dust, cigarette smoke, etc.).

Instructions for use

Use the dosing pipette provided.

Insert the pipette into the bottle and pull the plunger out as far as the black line. This is one 5 ml dose. Place the tip of the pipette in the baby's mouth. Gently press down on the plunger so that the syrup slowly enters the mouth and the baby can swallow it.

Dose

Babies from 6 to 12 months: 1 x 5 ml, max. 3 to 4 times a day.

Children from 1 to 6 years: 2 x 5 ml, max. 3 to 4 times a day.

Children from 6 to 12 years: 1 tablespoon (15 ml), max. 3 to 4 times a day.

Adults and children from 12 years: 1 tablespoon (15 ml), max. 4 to 6 times a day.

Do not exceed the above dose. If you forget to give a dose, do not double the dose.

The pipette must be cleaned with water after each use.

Do not clean the pipette in the dishwasher, do not use heat sterilisation and do not place the pipette in the microwave.

Warnings

Do not use with:

- babies younger than 6 months.
- allergy to one of the ingredients of this syrup
- hereditary fructose intolerance

It is advisable to consult your doctor for babies from 6 to 12 months.

You should also consult your doctor if dry cough is accompanied by severe symptoms (difficulty in breathing, fever lasting more than 3 days, repeated vomiting, feeding problems, diarrhea, changes in behavior or sleep disorders).

Avoid contact with the eyes and nose. In the event of accidental contact with the nose, eyes or ears of your baby / child, you need to rinse the syrup, preferably with physiological serum.

Keep the pipette out of the reach of children.

Ingredients

Glycerol, purified water, maltitol syrup, citric acid monohydrate, strawberry flavour, potassium sorbate, sodium benzoate

Shelf life

Store below 25 °C.

KEEP OUT OF THE SIGHT AND REACH OF CHILDREN.

Do not use Maga syrup after the date on the box and the label.

After opening, the syrup must be used within 3 months.

Presentation and delivery

180 ml bottle with 5 ml dosing pipette.

Free delivery.

Date of CE marking: 2016

Date of latest revision of information leaflet: April 2020.



Manufacturer:

Laboratoria Qualipharm, Rijksweg 9, 2880 Bornem, Belgium



Manufacturer (the pipette):

Bormioli Pharma S.p.A. Via Tortona 48, 27055 Rivanazzano (PV) ITALY



0373

Authorised representative of a manufacturer, importer and distributor:

EUPHARM phc d.o.o. Bilećka 37,

11000 Beograd, Republika Srbija

www.eupharm.com

www.drsmart.rs

Registration number: 515-02-03192-22-003

Catalogue number: 62950041